

CAPITULO 6

“BIBLIOGRAFÍA”



CAPITULO 6: BIBLIOGRAFÍA

AMAP. 2000. PCB in the Russian Federation: Inventory and Proposal for Priority Remedial Actions. Executive Summary. Arctic Monitoring and Assessment Programme, OSLO and Centre for International Projects, Moscow. <http://www.amap.no/ol-docs/pcb-es.pdf>

ATSDR (2000). Toxicological Profile for Polychlorinated Biphenyls. Agency for Toxic Substances and Disease Registry.

Arcaro, K. F. et al 1999. Antiestrogenicity of environmental polycyclic aromatic hydrocarbons in human breast cancer cells. *Toxicology*, 133: 115-127.)

Barra R., Quiroz R., Mendoza G., Urrutia R., K.Pozo and S. Focardi (2002a) Polychlorinated byphenils in Perumytilus purpuratus (Lamarck 1819) Along the Chilean coast reflects latitudinal concentration gradients. *Organohalogen compounds* 58:481-484.

Barra R., Urrutia R., Cisterna M., Muñoz P., Quiroz R., Pozo K. y S. Focardi (2002b). Bifenilos Policlorados en sedimento de lagos Chilenos. VI Encuentro de Química Analítica y Ambiental. Univ. Austral.

Barra R., Cisternas M., Suarez C., Araneda A., Piñones O. y P. Popp (2004a). PCBs and HCHs in a Saltmarsh Sediment Record from South Central Chile: Use of Tsunami Signatures and 137 Cs Fallout as Temporal Markers. *Chemosphere* (en prensa).

Barra R., Pozo K., Muñoz P., Araneda A., Urrutia R. y S. Focardi (2004b). PCBs in a Dated Sediment Core of a High Altitude Lake in the Chilean Altiplano: The Chungara Lake. *Fresenius Environmental Bulletin* (en prensa)

Blais J., Froese K.; Kimpe L, Muir D., Backus S., Comba M. and D. Schindler. Assessment and characterization of polychlorinated near a hazardous waste incinerator: analysis and sediments. *Environ. Tox. Chem.* 22 (1) 126-133.

Breivik K., Sweetman A., Pacyna J., Jones K. (2002). Towards a global historical inventory for selected PCB congeners- a mass balance approach 1. Global production and consumption. *The Sci. Total Environ.* 290: 181-198.

Biessmann A. Effects of polychlorinated biphenyls on gonads, sex hormone balance and reproduction processes of Japanese quail *Coturnix coturnix japonica* after ingestion during sexual maturation. *ENVIRON POLLUT SER A ECOL BIOL*; 27 (1). 1982. 15-30

Borghini F., Grimalt J., Barra R., Sánchez-Hernandez J. C., Torres Gracia C. and S. Focardi (2005). Temperature dependence of the organochlorine compound distribution in soils and sediments in the Andes Mountains. *Environmental Pollution*.

Carpenter, D. O. (1998). Polychlorinated Biphenyls and Human Health. *International Journal of Occupational Medicine and Environmental Health*, 11(4): 291-303

CEC. 1996. Status of PCB management in North America. Commission for Environmental Cooperation, Montreal. http://www.cec.org/pubs_info_resources/publications/pdfs/english/pcbe.pdf

CENMA (2001). Caracterización de los Bifenilos Policlorados (PCBs) en la atmósfera urbana de la Región Metropolitana de Chile.

CONAMA IV Región (2000). Diagnóstico Regional sobre presencia y situación de contaminantes orgánicos persistentes (COPs) como existencias obsoletas, Región de Coquimbo.

CONAMA (2004). Manual de Chile sobre el manejo de Bifenilos Policlorados . 114 pp.

Convenio de Basilea. Sobre el control de los movimientos transfronterizos de los desechos peligrosos y su eliminación (1989)

Convenio de Estocolmo. Sobre contaminantes orgánicos persistentes (COPs) (2001)

De Guise S, Martineau D, Beland P, Fournier M. Possible mechanisms of action of environmental contaminants on St. Lawrence beluga whales (*Delphinapterus leucas*). *Environ Health Perspect.* 1995 May;103 Suppl 4:73-7.

De Voogt P, Brinkman UATH: Production, properties and usage of polychlorinated biphenyls. In: Kimbrough RD, Jensen AA, editors, *Halogenated biphenyls, Terphenyls, naphtalenes, dibenzodioxins and related products. Topics in environmental health.* ISBN 0-444-81029-3 Elsevier, 1989:3-45

EPA (2000). Hudson River PCBs Reassessment RI/FS Phase 3 Report: Feasibility Study. U. S. Environmental Protection Agency, and U. S. Army Corps of Engineers. Online at <http://www.epa.gov/hudson/>

Focardi, S., C. Fossi, C. Leonzio, S. Corsolini and O. Parra. 1996. Persistent organochlorine residues in fish and water birds from the Bio Bio river, Chile. *Environ. Monitor. Assess.*, 43:73-92.

FONDECYT "Búsqueda de evidencias del transporte atmosférico a gran escala de contaminantes: acumulación de Contaminantes Orgánicos Persistentes (COPs) en sedimentos de lagos remotos de Chile" <http://www.conicyt.cl/bases/fondecyt/proyectos/01/2001/1010640.html>

Fuentealba. M. (1997). Pesticidas Organoclorados y Bifenilos Policlorados en *Trachurus murphyi* en la Zona Centro Sur de Chile. Bol Soc. Biol. Concepción 68:39-46.

Geyer H., Freitag D., Korte F. (1984): Polychlorinated biphenyls (PCBs) in the marine environment, particularly in the Mediterranean. Ecotox. Environ. Saf., 8, 129-151
International Mussel Watch, 1995. Initial implementation phase. Final report. Farrington, J. W. & B.W. Tripp, Eds. Woods Hole, Massachusetts. 63 pp.

Gladen B, Rogan W and Hardy P. (1988). Development after exposure to Polychlorinated biphenyls and Dichlorodiphenyl dichloroethene transplacentally and through human milk. J Pediatr: 113: 991-995.

IFCS. 2001. Framework for the management of PCB. Contribution to the IFCS PCB Strategy Group, Intergovernmental Forum on Chemical Safety. http://www.who.int/ifcs/pcb/pcb_framework.pdf

Gray, L. E. et al 1995. Functional Developmental Toxicity of Low Doses of 2,3,7,8-Tetrachlorodibenzo-p-Dioxin and a Dioxin-Like PCB (169) in Long Evans Rats and Syrian Hamsters: Reproductive, Behavioral and Thermoregulatory Alterations. Organohalogen Compounds, 25: 33.)

Helle E, Olsson M, Jensen S. 1976. Seals in the Baltic Sea and environmental poisons. FAUNA FLORA (STOCKH); 71 (2). 41-48

Jacobson, J. L. and Jacobson, S. W. (1996). Intellectual Impairment in Children Exposed to Polychlorinated Biphenyls in Utero. New England Journal of Medicine, 335(11): 783-789.

Johnson, B. L. et al (1999). Public Health Implications of Exposure to Polychlorinated Biphenyls (PCBs). Agency for Toxic Substances and Disease Registry. Online at <http://www.atsdr.cdc.gov/DT/pcb007.html>

Jiang K., Li L., Chen Y. and J. Jin. Determination of PCDD/Fs and dioxin-like PCBs in Chinese commercial PCBs and emissions from testing PCB incinerator. Chemosphere 1997; 34: 941-950

Kannan.K, Reusch T.B.H., Schultz-Bull D.E., Petrick G., Duinker J.C. (1995): Chlorobiphenyls: Model compounds for metabolism in food chain organisms and their potential use as ecotoxicological stress indicators by application of the metabolic slope concept. Environ. Sci. Technol., 29, 1851-1859.

Mandalakis M. and Stephanou E. (2002). Polychlorinated Biphenyls Associated with fine particles (PM_{2.5}) in the urban environment of Chile: Concentrations levels, and Sampling volatilization losses. Environ. Tox. and Chem. 21 (11) 2270-2275.

Mariottini, M.; S Aurigi and Focardi, S. (2000). Congener profile and toxicity assesment of polychlorinated biphenyls in human adipose tissue of Italians and Chileans. *Microchemical Journal* 67 63-71 pp.

Matta Mb, Cairncross C, Kocan Rm. 1998. Possible effects of polychlorinated biphenyls on sex determination in rainbow trout. *Environmental Toxicology And Chemistry*; 17 (1).. 26-29.

Mendola, P. et al, 1997. Consumption of PCB-contaminated Freshwater Fish and Shortened Menstrual Cycle Length. *American Journal of Epidemiology*, 145(11): 955.

Meijer, S. N.; Ockenden, W. A.; Sweetman, A.; Breivik, K.; Grimalt, J. O.; Jones, K. C. (2003). Global Distribution and Budget of PCBs and HCB in Background Surface Soils: Implications for Sources and Environmental Processes *Environ. Sci. Technol* 37(4); 667-672.

Muñoz, J and Becker, P. (1999). The Kelp Gull as bioindicator of Environmental Chemicals in the Magellan Region. A comparison with other coastal sites in Chile. *Sci. Marina* 63 (suppl1) 495-502.

Muñoz J., Becker P, Sommer U, Pacheco P, Schlatter R. (2003). Seabird eggs as bioindicators of chemical contamination in Chile. *Environmental Pollution* 126,123-137.

Notarianni V., Calliera M., Tremolada P., Finizio A., Vighi M. (1998). PCBs distribution in soil and vegetation from different areas in Northern Italy. *Chemosphere* 37, 2839-2845.

Orn S, Andersson PL, Forlin L, Tysklind M, Norrgren L 1998. The impact on reproduction of an orally administered mixture of selected PCBs in zebrafish (*Danio rerio*). *Arch Environ Contam Toxicol* Jul;35(1):52-7

Parra O., Valdovinos C., Figueroa and Acuña A. (1998). Bio-Bio river water quality program phase II 1995-1997. Administrative council of Bio-bio water quality monitoring program (ed) Concepción, Chile, pp 41.

Powell DC, Aulerich RJ, Stromborg KL, Bursian SJ 1996. Effects of 3,3',4,4'-tetrachlorobiphenyl, 2,3,3',4,4'-pentachlorobiphenyl, and 3,3',4,4',5-pentachlorobiphenyl on the developing chicken embryo when injected prior to incubation. *J Toxicol Environ Health* Oct 25;49(3):319-38

Rawn, D., Lockhart, L., Wilkinson, P, Savoi, D., Bruno, G., Muir, D., 2002. Historical contamination of Yukon Lake sediments by PCBs and organochlorine pesticides: influence of local sources and watershed characteristics. *Sci. Total Environ* 280: 17-37

Rose, N., Rippey, B., 2002. The historical record of PAH, PCB trace metal and fly-ash particle deposition at a remote lake in north-west Scotland. *Environ. Poll* 117: 121-132

Ruby S, Mendoza LT, Fournier M, Brousseau P, Degas V. 2003. Reproductive system impairment of mice fed diets containing beluga whale blubber from the St Lawrence estuary and arctic populations. *J Toxicol Environ Health A*. Jun 13;66(11):1073-85.

Sangalang GB, Freeman HC, Crowell R1 981. Testicular abnormalities in cod (*Gadus morhua*) fed Aroclor 1254. *Arch Environ Contam Toxicol*; VOL 10, ISS 5, , P617-26

Schell, L. M. et al 2000. Polychlorinated biphenyls and thyroid function in adolescents of the Mohawk Nation at Akwesasne. In *Proceedings of the Ninth International Conference*, Turin, Italy.

Schlosserová L. 'Kontrolle ausgewählter Böden in der Tschechischen und Slowakischen Republik auf ihre Kontamination mit chlorierten Kohlenwassertstoffen (In German). In: Heinisch E, Kettrup A, Wenzel-Klein S. editors. *Schadstoffatlas Osteuropa*. ISBN 3-609-69540-4 Germany: ecomed verlagsgesellschaft AG & Co, 1994: 54-59.

Stewart, P. et al 2000. Prenatal PCB exposure and neonatal behavioral assessment scale (NBAS) performance. *Neurotoxicology and Teratology*, 22: 21-29.)

Subramanian A, Tanabe S, Tatsukawa R, Saito S, Miyazaki N. Reduction in the testosterone levels by PCBs and DDE in Dall's porpoises of northwestern North Pacific. *MAR POLLUT BULL*; 18 (12). 1987. 643-646.

Tatsukawa R. PCB pollution of the Japanese environment. In: Higuchi K, editor. *PCB poisoning and pollution*. ISBN 0-12-347850-2 Kodansha Ltda. Tokyo: Academic Press, 1976: 147-179.

Tamayo C., R., M. N. Matus, S. C. Montes & V. R. Cristi. 1994. Polychlorinated biphenyls (PCBs) determination in human milk samples collected in three provinces of the tenth region of Chile 1990. (Facultad de Ciencias Veterinarias, Universidad Austral de Chile, Valdivia, Chile). *Rev. Med. Chile* 122 (7), 746-53.

UNEP. 1999. National inventories of persistent organic pollutants, selected examples and possible models. Preliminary report. Note by the Secretariat, UNEP Chemical, Geneva. (UNEP/POPS/INC.3/INF/1). <http://www.chem.unep.ch/sc/documents/meetings/inc3/infenglish/inf3-1/inc3-1.pdf>

UNEP. 1999. Guidelines for the Identification of PCBs and Materials Containing PCBs. UNEP Chemical, Geneva. Internet:<http://www.chem.unep.ch/irptc/Publications/pcb1d1.pdf>;

UNEP (2002a). Regional Assessment of Persistent Toxic Substances. Arctic Regional Report. UNEP Chemicals, Geneva, 145 pp.

UNEP (2002b). Regional Assessment of Persistent Toxic Substances. Antarctica Regional Report. UNEP Chemicals, Geneva, 76 pp.

UNEP (2002c). Regional Assessment of Persistent Toxic Substances. North America Regional Report. UNEP Chemicals, Geneva, 147 pp.

UNEP (2002d). Regional Assessment of Persistent Toxic Substances. Europe Regional Report. UNEP Chemicals, Geneva, 141 pp.

UNEP (2002e). Regional Assessment of Persistent Toxic Substances. Mediterranean Regional Report. UNEP Chemicals, Geneva, 148 pp.

UNEP (2002f). Regional Assessment of Persistent Toxic Substances. Sub-Saharan Africa Regional Report. UNEP Chemicals, Geneva, 118 pp.

UNEP (2002g). Regional Assessment of Persistent Toxic Substances. Indian Ocean Regional Report. UNEP Chemicals, Geneva, 93 pp.

UNEP (2002h). Regional Assessment of Persistent Toxic Substances. Central and Northeast Asia Regional Report. UNEP Chemicals, Geneva, 115 pp.

UNEP (2002i). Regional Assessment of Persistent Toxic Substances. Southeast Asia and South Pacific Regional Report. UNEP Chemicals, Geneva, 126 pp.

UNEP (2002j). Regional Assessment of Persistent Toxic Substances. Pacific Islands Regional Report. UNEP Chemicals, Geneva, 54 pp.

UNEP (2002k). Regional Assessment of Persistent Toxic Substances. Central America and the Caribbean Regional Report. UNEP Chemicals, Geneva, 133 pp.

UNEP (2002l). Regional Assessment of Persistent Toxic Substances. Eastern and Western South America Regional Report. UNEP Chemicals, Geneva, 90 pp.

UNEP (2003). Regional Assessment of Persistent Toxic Substances. Global Report. UNEP Chemicals, Geneva, 207 pp.

UNEP (2003). UNEP Workshop to develop a Global POPs Monitoring Programme to Support the Effectiveness Evaluation of the Stockholm Convention, Geneva, Switzerland, 24-27 March. 257 pp.

UNEP (2003). Global Monitoring Assessment Program. Workshop to develop a global assessment. Geneva March 2003.

Wade TI, Chambers L, Gardinali Pr, Sericano JI, Jackson TJ, Tarpley RJ, Suydam R. 1997. Toxaphene, PCB, DDT, and chlordane analyses of Beluga Whale blubber. *CHEMOSPHERE*; 34 (5-7).. 1351-1357.

Wania, F. D. Mackay (1993). Global fractionation and cold condensation of low volatility organochlorine compounds in polar regions. *Ambio* 22: 10-18.

Walker MK, Peterson RE. Toxicity of polychlorinated dibenzo-p-dioxins, dibenzofurans, and biphenyls during early development in fish. *Chemically-Induced Alterations in Sexual and Functional Development: the Wildlife/ Human Connection (Advances in Modern Environmental Toxicology; v.21)* 1992;:195-202.

Weinand-Harer, A. et al 1997. Behavioral effects of maternal exposure to an ortho-chlorinated or a coplanar PCB congener in rats. *Environmental Toxicology and Pharmacology*, 3: 97-103.)

Weisglas-Kuperus, N. et al 2000. Immunologic Effects of background Exposure to Polychlorinated Biphenyls and Dioxins in Dutch Preschool Children. *Environmental Health Perspectives*, 108(12): 1203)