

ANEXO 3

“REFERENCIAS MÉTODOS ANALÍTICOS PARA LOS DOCE COPS”



ANEXO 3: REFERENCIAS “MÉTODOS ANALÍTICOS PARA LOS DOCE COPs”

Abraham K, Hille A, Ende M, et al. 1994. Intake and fecal excretion of PCDDs, PCDFs, HCB and PCBs, (138,153,180) in a breast-fed and a formula-fed infant. *Chemosphere* 29(9-11):2279-2286.

Abraham K, Wiesmueller T, Brunner H, et al. 1989a. Elimination of various polychlorinated dibenzo-p-dioxins and dibenzofurans (PCDDs and PCDFs) in rat feces. *Arch Toxicol* 63:75-78.

Adeshina F, Todd EL. 1990. Organochlorine compounds in human adipose tissue from north Texas. *J Toxicol Environ Health* 29:147-156.

Aguilar C, Penalver A, Pocurull E, et al. 1999. Optimization of solid-phase microextraction conditions using a response surface methodology to determine organochlorine pesticides in water by gas chromatography and electron-capture detection. *J Chromatogr A* 844:425-432

Alawi MA, Ammari N, Al-Shuraiqi Y. 1992. Organochlorine pesticide contaminations in human milk samples from women living in Amman, Jordan. *Arch Environ Contam Toxicol* 23:235-239.

Aldrich FD, Holmes JH. 1969. Acute chlordane intoxication in a child: Case report with toxicological data. *Environ Health* 19:129-132.

Alford-Stevens AL, Bellar TA, Eichelberger JW, et al. 1986. Accuracy and precision of determinations of chlorinated pesticides and polychlorinated biphenyls with automatic interpretation of mass spectrometric data. *Anal Chem* 58(9):2022-2029.

Alford-Stevens AL, Eichelberger JW, Budde WL. 1988. Multilaboratory study of automated determinations of polychlorinated biphenyls and chlorinated pesticides in water, soil, and sediment by gas chromatography/mass spectrometry. *Environmental Science and Technology* 22:304-312.

Alley EG, Lu G. 1995. Analysis of polychlorinated biphenyls in fatty biological matrixes by on-line supercritical fluid extraction and supercritical fluid cleanup. *J AOAC Int* 78(4):1051-1054.

Andrews P, Newsome WH, Boyle M, et al. 1993. High resolution selective monitoring GC-MS determination of toxaphene in Great Lakes fish. *Chemosphere* 27(10):1865-1872.

AOAC. 1990. Association of Official Analytical Chemists. Agricultural chemicals; contaminants; drugs. Pesticides and industrial chemical residues. 1:274-280.

APHA. 1989a. Method 6410B. Liquid-liquid extraction gas chromatographic/mass spectrometric method. In: Clesceri LS, Greenberg AE, Trussels RR eds. *Standard methods for the examination of water and wastewater*. 17th ed. American Public Health Association, American Water Works Association, Water Pollution Control Federation 6-113 - 6-136.

APHA. 1989b. Method 6630c. Liquid-liquid extraction gas chromatographic method II. In: Clesceri LS, Greenberg AE, Trussels RR eds. Standard methods for the examination of water and wastewater. 17th ed. American Public Health Association, American Water Works Association, Water Pollution Control Federation 6-170 - 6-181.

ASTM. 1988. Method D 3086-85. Standard test method for organochlorine pesticides in water. 1988 annual book of ASTM standards. Vol. 11.02: Water and environmental technology. American Society for Testing and Materials, Philadelphia, PA 163-178.

ASTM. 1991. Standard practice for sampling and analysis of pesticides and polychlorinated biphenyls in indoor atmospheres. American Society for Testing and Materials. ASTM designation D 4861-91. p366-379.

Ault JA, Spurgeon TE, Gillard DS, et al. 1985. Multiresidue gas chromatographic method for determining organochlorine pesticides in meats validation study for swine and beef fats. J Assoc Off Anal Chem 68:941-944.

Ault JA, Spurgeon TE. 1984. Multiresidue gas chromatographic method for determining organochlorine pesticides in poultry fat: Collaborative study. J Assoc Off Anal Chem 67:284-289.

Baldwin MK, Hutson DH. 1980. Analysis of human urine for a metabolite of endrin by chemical oxidation and gas-liquid chromatography as an indicator of exposure to endrin. Analyst 105:60-65.

Barcarolo R, Tealdo E, Tutta C. 1988. Multiresidue determination of organochlorine and triazine pesticides in homogenized milk. J High Resolution Chromatog Chromatog Comm 11(7):539-541.

Barcarolo R, Tealdo E, Tutta C. 1988. Multiresidue determination of organochlorine and triazine pesticides in homogenized milk. J High Resolution Chromatography & Chromatography Comm 11:10746-10748.

Barquet A, Morgade C, Pfaffenberger CD. 1981. Determination of organochlorine pesticides and metabolites in drinking water, human blood serum, and adipose tissue. J Toxicol Environ Health 7:469-479.

Barrie L, Bidleman T, Dougherty D, et al. 1993. Atmospheric toxaphene in the high arctic. Chemosphere 27(10):2037-2046.

Behzadi H, Lalancette RA. 1991. Extending of the use of USEPA method 1625 for the analysis of 4,4'-DDT, 4,4'-DDD, and 4,4'-DDE. Microchem J 44:122-129.

Bidleman T, Matthews J, Olney C, et al. 1978. Separation of polychlorinated biphenyls, chlordane, and p,p'-DDD from toxaphene by silicic acid column chromatography. J Assoc Off Anal Chem 61(4):820-828.

Bigley WS, Plapp FW Jr, Hanna RL, et al. 1981. Effect of toxaphene, camphene and cedar oil on methyl parathion residues on cotton. Bull Environ Contam Toxicol 27:90-94.

Billings WN, Bidleman TF. 1980. Field comparison of polyurethane foam and tenax-GC resin for high-volume air sampling of chlorinated hydrocarbons. *Environmental Science and Technology* 14:679-683.

Bingham AG, Edmunds CJ, Graham BWL, et al. 1989. Determination of PCDDs and PCDFs in car exhaust. *Chemosphere* 19:669-673.

Bobbie BA, Clement RE, Taguchi VY. 1989. Determination of chlorinated dibenzo-p-dioxins (CDD) and dibenzofurans (CDF) in environmental samples by tandem mass spectrometry. *Chemosphere* 18:155-162.

Bong RL. 1975. Determination of hexachlorobenzene and mirex in fatty products. *J AOAC* 58(3):557-561.

Bong RL. 1977. Collaborative study of the recovery of hexachlorobenzene and mirex in butterfat and fish. *J Assoc Off Anal Chem* 60(1):229-232.

Boshoff PR, Pretorius V. 1979. Determination of toxaphene in milk, butter and meat. *Bull Environ Contam Toxicol* 22:405-412.

Brás I, Santos L, Alves A. 2000. Monitoring organochlorine pesticides from landfill leachates by gas chromatography-electron-capture detection after solid-phase microextraction. *J Chromatogr A* 891:305-311.

Bristol DW, Crist HL, Lewis RG, et al. 1982. Chemical analysis of human blood for assessment of environmental exposure to semivolatile organochlorine chemical contaminants. *J Anal Toxicol* 6:1-7.

Brownlow CS, Que Hee SS. 1985. Comparison of solid sampling media for Aroclor 1254 vapor under dry and humid conditions. *Am Ind Hyg Assoc J* 46:421-426.

Brumley WC, Brownrigg CM, Grange AH. 1993. Determination of toxaphene in soil by electron-capture negative-ion mass-spectrometry after fractionation by high-performance gel-permeation chromatography. *J Chromatogr* 633: 177-183.

Burse VW, Head SL, Korver MP, et al. 1990. Determination of selected organochlorine pesticides and polychlorinated biphenyls in human serum. *J Anal Toxicol* 14:137-146.

Burse VW, Korver MP, Needham LL, et al. 1989. Gas chromatographic determination of polychlorinated biphenyls (as Aroclor 1254) in serum: Collaborative study. *J Assoc Off Anal Chem* 72(4):649-659.

Burse VW, Needham LL, Korver MP, et al. 1983. Gas-liquid chromatographic determination of polychlorinated biphenyls and a selected number of chlorinated hydrocarbons in serum. *J Assoc Off Anal Chem* 66(1):32-39.

Buser HR, Rappe C, Bergqvist P. 1985. Analysis of polychlorinated dibenzofurans, dioxins, and related compounds in environmental samples. *Environ Health Perspect* 60:293-302.

Bush B, Barnard EL. 1982. Determination of nonpolar chlorinated hydrocarbons and polychlorinated biphenyls in microsamples. *Anal Lett* 15(20):1643-1648.

Bush B, Snow J, Connor S, et al. 1983a. Mirex in human milk in upstate New York. *Arch Environ Contam Toxicol* 12(6):739-746.

Bush B, Snow JT, Connor S. 1983b. High resolution gas chromatographic analysis of nonpolar chlorinated hydrocarbons in human milk. *J Assoc Off Anal Chem* 66(2):248-255.

Cai Z, Giblin DE, Ramanujam VMS, et al. 1994. Mass profile monitoring in trace analysis: Identification of polychlorodibenzothiophenes in crab tissue collected from the Newark/Raritan Bay system. *Environ Sci Technol* 28(8):1535-1538.

Caille G, Plaa GL, Vezina M. 1987. Gas-liquid chromatographic determination of chlordecone and mirex in biological specimens. *J Toxicol Clin Exp* 7(1):21-29.

Cairns T, Siegmund EG, Froberg JE. 1981. Chemical ionization mass spectrometric examination of metabolized toxaphene from milk fat. *Biomed Mass Spectrom* 8:569-574.

Calamari D, Tremolada P, Guardo AD, et al. 1994. Chlorinated hydrocarbons in pine needles in Europe: Fingerprint for the past and recent use. *Environ Sci Technol* 28:429-434.

Carey AE, Wiersma GB, Tai H. 1976. Pesticide residues in urban soils from 14 United States cities, 1970. *Pestic Monit J* 10(2):54-60.

Chan CH, Bruce G, Harrison B. 1994. Wet deposition of organochlorine pesticides and polychlorinated biphenyls to the great lakes. *J Great Lakes Res* 20(3):546-560.

Chang RR, Jarman WM, Hennings JA. 1993. Sample cleanup by solid phase extraction for the ultratrace determination of polychlorinated dibenzo-p-dioxins and dibenzofurans in biological samples. *Anal Chem* 65(18):2420-2427.

Chang RR, Jarman WM, King CC, et al. 1990. Bioaccumulation of PCDDs and PCDFs in food animals. III. A rapid cleanup of biological material using reverse-phase adsorbent columns. *Chemosphere* 20:881-886.

Chau ASY, Babjak LJ. 1979. Column chromatographic determination of mirex, photomirex, and polychlorinated biphenyls in lake sediments. *J Assoc Off Anal Chem* 62(1):107-113.

Clesceri LS, Greenberg AE, Eaton AD, et al. (eds.). 1998a. Method 6630C: Liquid-liquid extraction gas chromatographic method II. In: Standard methods for examination of water and wastewater. 20th ed. Washington, DC: American Public Health Association, 6.97-6.104.

Clesceri LS, Greenberg AE, Eaton AD, et al. (eds.). 1998b. Method 6410B: Liquid-liquid extraction gas chromatographic/mass spectrometric method. In: Standard methods for examination of water and wastewater. 20th ed. Washington, DC: American Public Health Association, 6.59-6.72.

Cooke M, Hatchel JA, Czuczwa JM, et al. 1988. Analytical and sampling techniques for the determination of ultratrace quantities of polychlorinated dibenzofurans and dibenzodioxins. In: Proceedings 1987 EPRI PCB Seminar, Kansas City, MO, October 6-9, 1987. Palo Alto, CA: Electric Power Research Institute, 7-1-7-8.

Creaser CS, Al-Haddad A. 1989. Fractionation of polychlorinated biphenyl, polychlorinated dibenzo-p-dioxins and polychlorinated dibenzofurans on porous graphitic carbon. *Anal Chem* 61:1300-1302.

Crist HL, Harless RL, Moseman RF, et al. 1980. Application of dehydrochlorination to the determination of toxaphene in soil and identification of the major gas chromatographic peak. *Bull Environ Contam Toxicol* 24:231-237.

Cruz I, Wells DE, Mat-r IL. 1993. Determination of organochlorines in sea water: An assessment. *Analytica Chimica Acta* 283:280-286.

Czuczwa J, Katona V, Pitts G, et al. 1989. Analysis of fog samples for PCDD and PCDF. *Chemosphere* 18:847-850.

Davies K. 1988. Concentrations and dietary intake of selected organochlorines, including PCBs, PCDDs and PCDFs in fresh food composites grown in Ontario, Canada. *Chemosphere* 17(2):263-276.

De la Riva C, Anadon A. 1991. Organochlorine pesticides in cow's milk from agricultural region in northwestern Spain. *Bull Environ Contam Toxicol* 46:527-533.

Di Muccio A, Rizzica M, Ausili A, et al. 1988. Selective, on-column extraction of organochlorine pesticide residues from milk. *J Chromatogr* 456(1):143-148.

Di Muccio A, Santilio A, Dommarco R, et al. 1990. Behaviour of 23 persistent organochlorine compounds during sulphuric acid clean-up on a solid-matrix column. *J Chromatogr* 513:333-337.

Djordjevic MV, Hoffmann D, Fan J, et al. 1994. Assessment of chlorinated pesticides and polychlorinated biphenyls in adipose breast tissue using a supercritical fluid extraction method. *Carcinogenesis* 15(11):2581-2585.

Donnelly JR, Vonnahme TL, Hedin CM, et al. 1986. Evaluation of RCRA method 8280 for analysis of dioxins and dibenzofurans. In: Rappe C, Choudhary G, Keith LH, eds. Chlorinated dioxins and dibenzofurans in perspective. Chelsea, MI: Lewis Publishers, Inc., 399-435.

Driscoll MS, Hassett JP, Fish CL, et al. 1991. Extraction efficiencies of organochlorine compounds from Niagara River water. *Environ Sci Technol* 25(8):1432-1439.

Durrell GS, Sauer TC. 1990. Simultaneous dual-column, dual-detector gas chromatographic determination of chlorinated pesticides and polychlorinated biphenyls in environmental samples. *Anal Chem* 62(17):1867-1871.

Eichelberger JW, Kerns EH, Olynx P, et al. 1983. Precision and accuracy in the determination of organics in water by fused silica capillary column gas chromatography/mass spectrometry and packed column gas chromatography/mass spectrometry. *Anal Chem* 55 : 1471 - 1479.

Eitzer BD, Chevalier A. 1999. Landscape care pesticide residues in residential drinking water wells. *Bull Environ Contam Toxicol* 62:420-427.

EMMI. 1997. Environmental Monitoring Methods Index. Version 1.1. PC# 4082. Rockville, MD: Government Institutes. U.S. Environmental Protection Agency.

EMMI. 1999. EPA environmental monitoring methods index: Detail analyte. Version I. PC no. 4082. Rockland, MD: Government Institutes (1997).

EPA 1986d. Test methods for evaluating solid waste. 3rd ed. Washington, DC: U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response. SW-846, 8080-I-8080-27, 8250-I-8250-32.

EPA 1998j. Method 8081B: Organochlorine pesticides by gas chromatography. U.S. Environmental Protection Agency.

EPA. 1976b. Analysis and GC-MS characterization of toxaphene in fish and water. U.S. Environmental Protection Agency, Office of Research and Development, Duluth, MN. (authors: Stalling DL, Huckins JN) EPA-600/3-76-076.

EPA. 1977. Manual of analytical methods for the analysis of pesticide residues in human and environmental samples. Revisions by the Assoc. of Official Analytical Chemists, U.S.EPA, Health Effects Research Laboratory, Environmental Toxicology Division, Research Triangle Park, NC.

EPA. 1980b. Analysis of pesticide residues in human and environmental samples: A compilation of methods selected for use in pesticide monitoring programs. U.S. Environmental Protection Agency. EPA 600/8-80-038.

EPA. 1980c. Analysis of pesticide residues in human and environmental samples: A compilation of methods selected for use in pesticide monitoring programs. Research Triangle Park, NC: U.S. Environmental Protection Agency, Health Effects Research laboratory, Environmental Toxicology Division. EPA-600/8-80-038.

EPA. 1980e. Manual of analytical methods for the analysis of pesticides in humans and environmental samples. U.S. Environmental Protection Agency. EPA-600/8-80-038.

EPA. 1982. Test Method-608: Organochlorine pesticides and PCB's. Cincinnati, OH: U.S. Environmental Protection Agency, environmental Monitoring and Support Laboratory.

EPA. 1982a. Test method: Organochlorine pesticides and PCBs - method 608. In: Longbottom JE, Lichtenberg JJ, eds. Test methods: Methods for organic chemical analysis of municipal and industrial wastewater. Cincinnati, OH: U.S. Environmental Protection Agency, Environmental Monitoring and Support Laboratory. EPA-600/4-82-057.

EPA. 1982a. Test methods for organic chemical analysis of municipal and industrial wastewater. Method No. S. 608 and 625. Environmental Monitoring and Support Laboratory, U.S. EPA, Cincinnati, OH.

EPA. 1982a. Test methods. Methods for organic chemical analysis of municipal and industrial wastewater. Cincinnati, OH: U.S Environmental Protection Agency, Office of Research and Development, Environmental Monitoring and Support Laboratory, 608-1 to 608-11; 625-1 to 625-12. EPA 600/4-82-057.

EPA. 1982b. Method development for determination of polychlorinated hydrocarbons in municipal sludge. U.S. Environmental Protection Agency, Office of Research and Development, Cincinnati, OH. (authors: Rodriguez CF, et al.) EPA-600/4-82-035.

EPA. 1982b. Test method: Base/neutrals and acids - method 625. In: Longbottom JE, Lichtenberg JJ, eds. Test methods: Methods for organic chemical analysis of municipal and industrial wastewater. Cincinnati, OH: U.S. Environmental Protection Agency, Environmental Monitoring and Support Laboratory. EPA-600/4-82-057.

EPA. 1984a. Method T04. Method for the determination of organochlorine pesticides and polychlorinated biphenyls in ambient air. Compendium of methods for the determination of toxic organic compounds in ambient air. U.S. Environmental Protection Agency, Quality Assurance Division, Environmental Monitoring Systems Laboratory, ORD, Research Triangle Park, NC. (authors: Winberry WT, et al.) EPA-600/4-84-041.

EPA. 1984c. EPA Method 608. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR Part 136.

EPA. 1986a. Test methods for evaluating solid wastes. SW-846, Method no. 8080, Volume 1B: Laboratory Manual: Physicachemical Methods, 3rd ed. Office of Solid Waste and Emergency Response, U.S. EPA, Washington, DC.

EPA. 1986b. Method 8250: Gas chromatography/mass spectrometry for semivolatile organics: Packed column technique. In: Test methods for evaluating solid waste. Volume IC: Laboratory manual: Physical/chemical methods. SW-846. 3rd ed. Washington, DC: U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response.

EPA. 1986b. Test method for evaluating solid waste. Physical/chemical methods. Washington, DC: U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response. 3rd Edition.

EPA. 1986c. Method 8270: Gas chromatography/mass spectrometry for semivolatile organics: Capillary column technique. In: Test methods for evaluating solid waste. Volume IC: Laboratory manual: Physical/chemical methods. SW-846. 3rd ed. Washington DC: U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response.

EPA. 1986f. Broad scan analysis of the FY82 national human adipose tissue survey specimens: Volume I - Executive summary. Washington, DC: U.S. Environmental Protection Agency, Office of Toxic Substances. EPA-560/5-86-035.

EPA. 1986f. Method 8080. Test methods for evaluating solid waste. Volume IB: Laboratory manual physical/chemical methods. SW 846, 3rd ed. U.S. Environmental Protection Agency, Office of Solid Waste, Washington, DC.

EPA. 1986j. Test methods for evaluating solid waste, 3rd ed. SW-846. Washington, DC: U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response. 8080-1-8080-27, 8250-1-8250-32.

EPA. 1986k. Test methods for evaluating solid waste (SW 846) volume IB: Laboratory manual physical chemical methods. Method 8280, the analysis of polychlorinated dibenzo-p-dioxins and polychlorinated dibenzofurans. U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response, Washington, DC.

EPA. 1987a. Health advisories for 16 pesticides (Toxaphene). U.S. Environmental Protection Agency, Office of Drinking Water, Washington, DC. PB87-200178.

EPA. 1987a. U.S. Environmental Protection Agency. Statement of work for organics analyses, multi-media, multiconcentration. U.S. EPA Contract Laboratory Program. Revised 8/87.

EPA. 1987a. U.S. EPA contract laboratory program. Statement of work for organic analysis. U.S. EPA, Washington, DC.

EPA. 1987f. Method 525.1: Determination of organic compounds in drinking water by liquid-solid extraction and capillary column gas chromatography/mass spectrometry. In: Methods for the determination of organic compounds in drinking water. Cincinnati, OH: U.S. Environmental Protection Agency, Environmental Monitoring Systems Laboratory.

EPA. 1988b. U.S. Environmental Protection Agency. Method 608: Organochlorine pesticides and PCBs. Code of Federal Regulations. 40 CFR 136, App. A, 354-374.

EPA. 1988c. Methods for the determination of organic compounds in drinking water. Cincinnati, OH: U.S. Environmental Protection Agency, Environmental Monitoring Systems Laboratory, Office of Research and Development. EPA-600/4-88-039.

EPA. 1988d. Compendium of methods for the determination of toxic organic compounds in ambient air. Research Triangle Park, NC: U.S. Environmental Protection Agency, Quality Assurance Division, Environmental Monitoring Systems Laboratory. EPA-600/4-89-017.

EPA. 1988e. Method 505: Method for the determination of organic compounds in drinking water. Washington, DC: U.S. Environmental Protection Agency. EPA 600/4-88/039.

EPA. 1988f. Method 508: Method for the determination of organic compounds in drinking water. Washington, DC: U.S. Environmental Protection Agency. EPA 600/4-88/039.

EPA. 1988g. Compendium on methods for the determination of toxic organic compounds in ambient air, Method TO-9, method for the determination of polychlorinated dibenzo-p-dioxins(PCDDs) in ambient air using high resolution gas chromatography high resolution mass spectrometry. (HRGC/HRMS). EPA 600/4/89/017.

EPA. 1988h. Analytical and sampling methods of the Nonoccupational Pesticide Exposure Study (NOPES). Washington, DC: U.S. Environmental Protection Agency. EPA 600/9-88-015.

EPA. 1989. Method 508. Determination of chlorinated pesticides in water by gas chromatography with an electron capture detector. In: Methods for the determination organic compounds in drinking water. Revision 3.0 (RL Graves). U.S. Environmental Protection Agency, Environmental Monitoring Station Lab, Cincinnati, OH. EPA-600/4-88-039.

EPA. 1989c. Methods for the determination of organic compounds in drinking water. Method No. 505, 508 and 508A. Cincinnati, OH: U.S. Environmental Protection Agency, Office of Research and Development, Environmental Monitoring Systems Laboratory.

EPA. 1992b. Method 1656: the determination of organo-halide pesticides in municipal and industrial wastewater. In: Methods for the determination of nonconventional pesticides in municipal and industrial wastewater. U.S. Environmental Protection Agency. EPA 821 RR-92-002. Pp 637-677.

EPA. 1992c. Method 617. The determination of organohalide pesticides and PCBs in municipal and industrial wastewater. In: Methods for the determination of nonconventional pesticides in municipal and industrial wastewater. EPA/821/R-92-002, 139-168.

EPA. 1992d. Method 1656. The determination of organo-halide pesticides in municipal and industrial wastewater. In: Methods for the determination of nonconventional pesticides in municipal and industrial wastewater. EPA/821/R-92-002. 637-677.

EPA. 1994a. Method 1613: tetra-through octa chlorinated dioxins and furans by isotope dilution HRGC/HRMS. United States Environmental Protection Agency, Office of Water, Engineering and Analysis Division, Washington DC. EPA 821-B-94-005.

EPA. 1994c. Polychlorinated dibenzodioxins (PCDDs) and polychlorinated dibenzofurans (PCDFs) by high-resolution gas chromatography/high-resolution mass spectrometry (HRGC/HRMS). Test methods for evaluating solid waste, method 8290. Environmental Protection Agency, Washington, DC.

EPA. 1994f. Method 8080A. Organochlorine pesticides and polychlorinated biphenyls by gas chromatography. U.S. Environmental Protection Agency.

EPA. 1995a. U. S. Environmental Protection Agency: Method 4040 soil screening for toxaphene by immunoassay. Test methods for evaluating solid waste, Vol IB: Laboratory manual physical/chemical methods. Revision 0, January 1995.

EPA. 1995c. Method 8082. Polychlorinated biphenyls (PCBs) by capillary column gas chromatography. U.S. Environmental Protection Agency.

EPA. 1995c. U. S. Environmental Protection Agency: Method 808 1 A: Chlorine pesticides by capillary column gas chromatography. Test methods for evaluating solid waste, Vol IB: Laboratory Manual Physical/Chemical methods. Revision 0, January 1995.

EPA. 1998k. Method 8270D: Semivolatile organic compounds by gas chromatography/mass spectrometry (GC/MS). Washington, DC: U.S. Environmental Protection Agency.

EPA. 1999k. Method 1668, Revision A: Chlorinated biphenyl congeners in water, soil, sediment, and tissue by HRGC/HRMS. U.S. Environmental Protection Agency, Office of Water. EPA-821-R-00-002.

Eschenroeder A, Jaeger RJ, Ospital JJ, et al. 1986. Health risk analysis of human exposures to soil amended with sewage sludge contaminated with polychlorinated dibenzodioxins and dibenzofurans. Vet Hum Toxicol 28: 435-442.

Fairless BJ, Bates DI, Hudson J, et al. 1987. Procedures used to measure the amount of 2,3,7,8-tetrachlorodibenzo-p-dioxin in the ambient air near a Superfund site cleanup operation. Environ Sci Technol 21:550-555.

FDA. 1994a. Food and Drug Administration: Method 302 for non-fatty food. Pesticide Analytical Manual, Vol 1. Mutiresidue Methods, 3rd edition.

FDA. 1994b. Food and Drug Administration: Method 304 for fatty food. Pesticide Analytical Manual, Vol 1. Multiresidue Methods. 3rd edition.

Ferrario J, Byrne C, McDaniel D, et al. 1996. Determination of 2,3,7,8-chlorine substituted dibenzo-p-dioxins and furans at the part per trillion level in United States beef fat using high resolution gas chromatography/high resolution mass spectrometry. *Anal Chem* 68:647-652.

Ford WM, Hill EP. 1991. Organochlorine pesticides in soil sediments and aquatic animals in the upper Steele bayou watershed of Mississippi. *Arch Environ Contam* 20:161-167.

Först C, Stieglitz L, Zwick G. 1988. Isomer-specific determination of PCDD/PCDF in oil extracts from water leachate of a waste landfill. *Chemosphere* 17:1935-1944.

Fuoco R, Colombini MP, Samcova E. 1993. Individual determination of ortho and non-ortho substituted polychlorobiphenyls (PCBs) in sediments by high performance liquid chromatographic pre-separation and gas chromatography/ECD detection. *Chromatographia* 36:65-70.

Games ED. 1977. Determination of toxaphene by basic alcoholic hydrolysis and florisil separation *Bull Environ Contam Toxicol* 17:456-462.

Ganguly SK, Bhattacharyya J. 1973. Detection of small amounts of pesticides in human biological material by thin-layer chromatography. *Forensic Sci* 2:333-338.

Garrison AW, Pellizzari ED. 1987. Application of the master analytical scheme to polar organic compounds in drinking water. In: Suffet IH, Malaiyandi M, eds. *Organic pollutants in water: Sampling, analysis, and toxicity testing*. Advances in Chemistry Series No. 214. American Chemical Society.

Gibson JR, Ivie GW, Dorrough HW. 1972. Fate of mirex and its major photodecomposition product in rats. *J Agric Food Chem* 20: 1246-1248.

Glidden RM, Brown PJ, Sittig RA, et al. 1990. Determination of 2,3,7,8-tetra-chlorodibenzop-dioxin and 2,3,7,8-tetrachlorodibenzofuran in cow's milk. *Chemosphere* 20:1619-1624.

Gore RC, Hannah RW, Pattacini SC, et al. 1971. Infrared and ultraviolet spectra of seventy-six pesticides. *J Assoc Off Anal Chem* 54(5):1040-82.

Graham M, Hileman FD, Orth RG, et al. 1986. Chlorocarbons in adipose tissue from a Missouri population. *Chemosphere* 15:1595-1600.

Graupner AJ, Dunn CL. 1960. Determination of toxaphene by a spectrophotometric diphenylamine procedure. *J Agric Food Chem* 8:286-289.

Greizerstein HB, Gigliotti P, Vena J et al. 1997. Standardization of a method for the routine analysis of polychlorinated biphenyl congeners and selected pesticides in human serum and milk. *J Analytical Toxicol* 21:125-133.

Griffith FD, Blanke RV. 1974. Microcoulometric determination of organochlorine pesticides in human blood. *J Assoc Off Anal Chem* 57:595-603.

Grob K, Kalin I. 1991. Attempt for an on-line size exclusion chromatography-gas chromatography method for analyzing pesticide residues in foods. *J Agric Food Chem* 39:1950-1953.

Halt RL, Cruse S, Greer ES. 1986. Pesticide and polychlorinated biphenyl residues in human adipose tissue from Northeast Louisiana. *Bull Environ Contam Toxicol* 36(5):651-655.

Hardin ED, Hayes DA, Shore FL, et al. 1989. A comparison of analytical methods for the analysis of chlorinated dioxin/dibenzofuran in pyrolyzed transformer oil samples. *Chemosphere* 18:1093-1100.

Hargesheimer EE. 1984. Rapid determination of organochlorine pesticides and polychlorinated biphenyls, using selected ion monitoring mass spectrometry. *J Assoc Off Anal Chem* 67(6):1067-1075.

Harkov R. 1986. Semivolatile organic compound in the atmosphere: A review. *J Environ Sci Health* 409-433.

Harless RL, Lewis RG, McDaniel DD, et al. 1992. Evaluation of a sampling and analysis method for determination of polyhalogenated dibenzo-p-dioxins and dibenzofurans in ambient air. *Chemosphere* 25(7-10):1317-1322.

Head SL, Burse VW. 1987. Organochlorine recovery from small adipose samples with the universal trace residue extractor (Unitrex). *Bull Environ Contam Toxicol* 39:848-856.

Hellou J, Warren WG, Payne JF. 1993. Organochlorines including polychlorinated biphenyls in muscle, liver and ovaries of cod, *Gadus morhua*. *Arch Environ Contam Toxicol* 25:497-505.

Helrich K, (ed). 1990. Method 970.52: Organochlorine and organophosphorus pesticide residues: General multiresidue methods. In: *Official methods of analysis of the Association of Official Analytical Chemists*. 15th ed. Arlington, VA: Association of Official Analytical Chemists, Inc., 274-282.

Helrich K, ed. 1990. Pesticide and industrial chemical residues. In: Official Methods of Analysis of the Association of Official Analytical Chemists. Arlington, VA: Association of Official Analytical Chemists, Inc., 274-284.

Hernandez HF, Lopez BFJ, Medina EJ, et al. 1987. Sulfuric acid cleanup and potassium hydroxideethanol treatment for confirmation of organochlorine pesticides and polychlorinated biphenyls: Application to wastewater samples. *J Assoc Off Anal Chem* 70:727-733.

Hippelein M, Kaupp H, Doerr G, et al. 1993. Testing of a sampling system and analytical method for determination of semivolatile organic compounds in ambient air. *Chemosphere* 26(12):2255-2263.

Ho JS, Tang PH, Eichelberger JW, et al. 1995. Liquid solid disk extraction followed by SFE and GC-ion-trap MS for the determination of trace organic pollutants in water. *J Chromatogr Sci* 33(1):1-g.

Hogmire HW, Weaver JE, Brooks JL. 1990. Survey for pesticides in wells associated with apple and peach orchards in West Virginia. *Bull Environ Contam Toxicol* 44:81-86.

Hong C-S, Bush B, Xiao J. 1992a. Isolation and determination of mono-ortho and non-ortho substituted PCBs (coplanar PCBs) in human milk by HPLC porous graphitic carbon and GC-ECD. *Chemosphere* 24(4):465-473.

Hopper M, King JW. 1991. Enhanced supercritical fluid carbon dioxide extraction of pesticides from foods using pelletized diatomaceous earth. *J Assoc Off Anal Chem* 74(4):661-666.

Hopper ML. 1982. Automated gel permeation system for rapid separation of industrial chemicals and organophosphate and chlorinated pesticides from fats. *J Agric Food Chem* 30:1038-1041.

Hsu JP, Schattenberg HJ, Garza MM. 1991. Fast turnaround multiresidue screen for pesticides in produce. *J Assoc Off Anal Chem* 74:886-892.

Hsu JP, Wheeler Jr HG, Camann DE, et al. 1988. Analytical methods for detection of nonoccupational exposure to pesticides. *J Chromatogr Sci* 26:181-189.

Hunt DF, Shabanowitz J, Harvey TM, et al. 1985. Scheme for the direct analysis of organics in the environment by tandem mass spectrometry. *Anal Chem* 57:525-537.

Ismail RJ, Bonner FL. 1974. New, improved thin layer chromatography for polychlorinated biphenyls, toxaphene, and chlordane components. *J Assoc Off Anal Chem* 57:1026-1032.

Jamuzi G, Wakimoto T. 1991. Determination of chlorinated pinene originated from pulp mill. *Analytical Sciences* V.7 Supplement: 1177-I 180.

Janak K, Becker G, Colmsjo A, et al. 1998. Methyl sulfonyl polychlorinated biphenyls and 2,2-bis(4-chlorophenyl)-1,1-dichlorethene in gray seal tissues determined by gas chromatography with electron capture detection and atomic emission detection. *Environ Toxicol Chem* 17(6):1046-1055.

Jansson B, Andersson R, Asplund L, et al. 1991. Multiresidue method for the gas-chromatographic analysis of some polychlorinated and polybrominated pollutants in biological samples. *Fresenius J Anal Chem* 340:439-445.

Jansson B, Wideqvist U. 1983. Analysis of toxaphene (PCC) and chlordane in biological samples by NCI mass spectrometry. *Int J Environ Anal Chem* 13:309-321.

Jasinski JS. 1989. Multiresidue procedures for the determination of chlorinated dibenzodioxins and dibenzofurans in a variety of foods using capillary gas chromatography-electron-capture detection. *J Chromatogr* 478:349-367.

Kelly AG, Cruz I, Wells DE. 1993. Polychlorobiphenyls and persistent organochlorine pesticides in sea water at the pg l1 level. Sampling apparatus and analytical methodology. *Anal Chim Acta* 276(1):3-13.

Klein AK, Link JD, Ives NF. 1968. Isolation and purification of metabolites found in the urine of male rats fed aldrin and dieldrin. *J Assoc Anal Chem* 51:895-898.

Klemmer HW, Budy AM, Takahashi W. 1977. Human tissue distribution of cyclodiene pesticides—Hawaii 1964-1973. *Clin Toxicol* 11(1):71-82.

Kleopfer RD, Greenall RL, Viswanathan TS, et al. 1989. Determination of polychlorinated dibenzodioxins and dibenzofurans in environmental samples using high resolution mass spectrometry. *Chemosphere* 18:109-118.

Kobayashi H, Sato K, Matano O, et al. 1983. Determination of aldrin and dieldrin in soil by mass chromatography. *J Pest Sci* 8(1):105-110.

Korfmacher WA, Rushing LG, Nestorick DM, et al. 1985. Analysis of dust and surface swab samples for octachlorodibenzo-p-dioxin and heptachlorodibenzo-p-dioxins by fused silica capillary GC with EC detection. *Journal of High Resolution Chromatography and Chromatography Communications* 8:12-19.

Korver MP, Burse VW, Needham LL, et al. 1991. Determination of mirex in human blood serum containing polychlorinated biphenyls by using packed column gas chromatography. *J Assoc Off Anal Chem* 74(5):875-877.

Krause RT. 1973. Determination of several chlorinated pesticides by the AOAC multiresidue method with additional quantitation of perthane after dehydrochlorination: Collaborative study. *J Assoc Off Anal Chem* 56(3):721-727.

Krauthacker B, Alebic-Kolbah T, Buntic A, et al. 1980. DDT residues in samples of human milk, and in mothers' and cord blood serum, in a continental town in Croatia (Yugoslavia). *Int Arch Occup Environ Health* 46:267-273.

Kraut-Vass A, Thoma J. 1991. Performance of an extraction disk in synthetic organic chemical analysis using gas chromatograph-mass spectrometry. *J Chromatog* 538:233-240.

Kurtz D. 1977. Adsorption of PCB's and DDT's on membrane filters- A new analysis method. *Bull Environ Contam Toxicol* 17(4):391-398.

Kutz FW, Strassman SC, Stroup CR, et al. 1985. The human body burden of mirex in the southeastern United States. *J Toxicol Environ Health* 15(3-4):385-394.

Kutz FW, Yobs AR, Yang HSC. 1976. National pesticide monitoring programs. In: Lee RI, ed. *Air pollution from pesticide and agricultural processes*, 95-137.

Kuwata K, Yamashita Y, Nakashima S, et al. 1993. Practical method for monitoring polychlorinated dibenzo-p-dioxins and polychlorinated dibenzofurans in the atmosphere. *J Chromatogr* 643(1-2):25-33.

LaFleur L, Bosquet T, Ramage K, et al. 1990. Analysis of TCDD and TCDF on the ppq-level in milk and food sources. *Chemosphere* 20:1657-1662.

Lamparski LL, Nestrick TJ. 1989. Analytical methodology of the Dow Chemical Company for the determination of selected chlorinated dibenzo-p-dioxins and dibenzofurans in stack gas effluent matrices: Part 1: Full method for determination of tetra- through octa-chlorinated congener group total concentrations and 2378 substituted isomers. *Chemosphere* 19:1165-1178.

Langhorst ML, Nestrick TJ. 1979. Determination of chlorobenzenes in air and biological samples by gas chromatography with photoionization detection. *Anal Chem* 51:2018-2025.

Le Be1 GL, Williams DT, Benoit FM. 1991. Determination of chlorinated dibenzodioxins and dibenzofurans in selected paper products. *Chemosphere* 23:737-746.

Le Bel GL, Williams DT. 1986. Determination of halogenated contaminants in human adipose tissue. *J Assoc Off Anal Chem* 69:451-458.

Lee, H-B, Peart TE. 1994. Optimization of supercritical carbon dioxide extraction for polychlorinated biphenyls and chlorinated benzenes from sediments. *J Chromatogr* 663:87-95.

Leister DL, Baker JE. 1994. Atmospheric deposition of organic contaminants to the Chesapeake Bay. *Atmos Environ* 28(8):1499-1520.

Lewis RG, Bond AE, Fitz-Simons TR, et al. 1986. Monitoring for non-occupational exposure to pesticides in indoor and personal respiratory air. In: proceedings of the 79th Annual meeting of the air pollution control association, June 22-27, 1986. Minneapolis, MN: APCA, 86-37.4:1-15.

Lewis RG, Brown AR, Jackson MD. 1977. Evaluation of polyurethane foam for sampling of pesticides, polychlorinated biphenyls and polychlorinated naphthalenes in ambient air. *Anal Chem* 49(12):1668-1672.

Liao W, Joe T, Cusick WG. 1991. Multiresidue screening method for fresh fruit and vegetables with gas chromatographic/mass spectrometric detection. *J Assoc Off Anal Chem* 74(3):554-565.

Ligocki MP, Pankow JF. 1985. Assessment of adsorption/solvent extraction with polyurethane foam and adsorption/thermal desorption with Tenax-GC for the collection and analysis of ambient organic vapors. *Anal Chem* 57:1138-1144.

Ligon WV, May RJ. 1986. Determination of selected chlorodibenzofurans and chlorodibenzodioxins using two-dimensional gas chromatography- mass spectrometry. *Anal Chem* 58:558-561.

Lin JM, Que Hee SS. 1985. Optimization of perchlorination conditions for some representative polychlorinated biphenyls. *Anal Chem* 57:2130-2134.

Lin JM, Que Hee SS. 1987. Change in chromatogram patterns after volatilization of some Aroclors, and the associated quantitation problems. *Am Ind Hyg Assoc J* 48:599-607.

Lindstroem G, Rappe C. 1990. Analytical procedure operating on 0.1-l ppt level applied in toxicological studies on PCDFs/PCDDs. *Chemosphere* 20:851-856.

Lopez-Avila V, Wesselman R, Edge11 K. 1990. Gas chromatographic-electron capture detection method for determination of 29 organochlorine pesticides in finished drinking water: Collaborative study. *J Assoc Off Anal Chem* 73(2):276-286.

Luke MA, Froberg JE, Masumoto HT. 1975. Extraction and clean-up of organochlorine, organophosphate; organonitrogen, and hydrocarbon pesticides in produce for determination by gas-liquid chromatography. *J Assoc Off Anal Chem* 58:1020-1026.

Luotamo M, Iarvisalo J, Aitio A. 1985. Analysis of polychlorinated biphenyls (PCBs) in human serum. *Environ Health Perspect* 60:327-332.

MacCuaig RD. 1976. The occurrence of insecticides in the blood of staff of a locust control organization. *Bull Environ Contam Toxicol* 15:162-170.

Macleod KE, Hanisch RC, Lewis RG. 1982. Evaluation of gel permeation chromatography for clean up of human adipose tissue samples for gas chromatographic-mass spectrometric analysis of pesticides and other chemicals. *J Anal Toxicol* 6(1):38-40.

Maiorino RM, Whiting FM, Brown WH, et al. 1980. Quantitative determination of toxaphene in blood by florisil extraction and gas chromatography. *J Anal Toxicol* 4:192-198.

Manes J, Font G, Pica Y. 1993. Evaluation of a solid-phase extraction system for determining pesticide residues in milk. *J Chromatog* 642:195-204.

Manes J, Font G, Pico Y. 1993. Evaluation of a solid-phase extraction system for determining pesticide residues in milk. *J Chromatogr* 642:195-204.

Marquis PJ, Hacket M, Holland LG, et al. 1994. Analytical methods for a national study of chemical residues in fish. 1: Polychlorinated dibenzo-p-dioxins, dibenzofurans. *Chemosphere* 29(3):495-508.

Marsh JM. 1993. Assessment of nonpoint source pollution in stormwater runoff in Louisville, (Jefferson County) Kentucky, USA. *Arch Environ Contam Toxicol* 25:446-455.

McCurvin DMA, Schellenberg DH, Clement RE, et al. 1989. A comparison of tandem quadrupole mass spectrometry versus single quadrupole capabilities for the determination of chlorinated dioxins and dibenzofurans in environmental samples. *Chemosphere* 19:201-204.

McKinney J, Moore L, Prokopetz A, et al. 1984. Validated extraction and cleanup procedures for polychlorinated biphenyls and DDE in human body fluids and infant formula. *J Assoc Off Anal Chem* 67(1):122.

McMahon B, Burke J. 1978. Analytical behavior data for chemicals determined using AOAC multiresidue methodology for pesticide residues in foods. *J Assoc Off Anal Chem* 61(3):640-652.

Mes J, Arnold DL, Bryce F. 1995a. The elimination and estimated half-lives of specific polychlorinated biphenyl congeners from the blood of female monkeys after discontinuation of daily dosing with Aroclor® 1254. *Chemosphere* 30:789-800.

Mes J, Arnold DL, Bryce F. 1995b. Postmortem tissue levels of polychlorinated biphenyls in female Rhesus monkeys after more than six years of daily dosing with Aroclor® 1254 and in their non-dosed offspring. *Arch Environ Toxicol* 29:69-76.

Mes J, Davies DJ, Turton D. 1982. Polychlorinated biphenyl and other chlorinated hydrocarbon residues in adipose tissue of Canadians. *Bull Environ Contam Toxicol* 28:97-104.

Mes J, Doyle JA, Adam BR, et al. 1984. Polychlorinated biphenyls and organochlorine pesticides in milk and blood of Canadian women during lactation. *Arch Environ Contam Toxicol* 13:217-223.

Mes J. 1992. Organochlorine residues in human blood and biopsy fat and their relationship. *Bull Environ Contam Toxicol* 48(6):815-820.

Mes J. 1994. Temporal changes in some chlorinated hydrocarbon residue levels of Canadian breast milk and infant exposure. *Environ Pollut* 84(3):261-268.

Miskiewicz AG, Gibbs PJ. 1994. Organochlorine pesticides and hexachlorobenzene in tissues of fish and invertebrates caught near a sewage outfall. *Environ Pollut* 84:269-277.

Miyata H, Aozasa O, Ohta S, et al. 1993. Estimated daily intakes of PCDDs, PCDFs and non-ortho coplanar PCBs via drinking water in Japan. *Chemosphere* 26(8):1527-1536.

Mühlebach S, Wyss P, Bickel M. 1985. Comparative adipose tissue kinetics of thiopental, DDE and 2,4,5,2',4',5'-hexachlorobiphenyl in the rat. *Xenobiotica* 15(6):485-491.

Munch DJ, Maxey RA, Engel TM. 1990. Methods development and implementation for the National Pesticide Survey. *Environ Sci Technol* 24(10):1446-1451.

Murk AJ, Legler J, Denison MS, et al. 1996. Chemical-activated luciferase gene expression (calux): A novel in vitro bioassay for ah receptor active compounds in sediments and pore water. *Fundam Appl Toxicol* 33(1):149-60.

Mussalo-Rauhamaa H, Pyysalo H, Antervo K. 1993. Relation between the content of organochlorine compounds in Finnish human milk and characteristics of the mothers. *J Toxicol Environ Health* 25:1-19.

Mussalo-Rauhamaa H. 1991. Partitioning and levels of neutral organochlorine compounds in human serum, blood cells, and adipose and liver tissue. *Sci Total Environ* 103:159-175.

Muto H, Takizawa Y. 1989. Dioxins in cigarette smoke. *Arch Environ Health* 44:171-174.

Nachman G, Freal J, Barquet A, et al. 1972. A simplified method for analysis of DDT and DDE in blood for epidemiologic purposes. In: Davies J, Edmundson, W, eds. *Epidemiology of DDT*. Mount Kisco, NY: Futura Publishing Company, Inc., 133-139.

Nam KS, King JW. 1994. Coupled SFE/SFC/GC for the trace analysis of pesticide residues in fatty food samples. *Journal of High Resolution Chromatography* 17:577-582.

Needham LL, Burse VW, Price HA. 1981. Temperature-programmed gas chromatographic determination of polychlorinated and polybrominated biphenyls in serum. *J Assoc Off Anal Chem* 64:1131-1137.

Needham LL, Smrek AL, Head SL, et al. 1980. Column chromatography separation of polychlorinated biphenyls from dichlorodiphenyltrichloroethane and metabolites. *Anal Chem* 2:2227-2229.

Nestrick TJ, Lamparski LL, Cramm RH. 1991. Procedure and initial results for the determination of low parts-per-quadrillion levels of 2,3,7,8-tetrachlorodibenzo- p-dioxin and 2,3,7,8-tetrachlorodibenzofuran in polyethylene. *Chemosphere* 22:215-228.

Newman JW, Vedder J, Jarman WM, et al. 1994. A method for the determination of environmental contaminants in living marine mammals using microscale samples of blubber and blood. *Chemosphere* 28(10):1795-1805.

NIOSH. 1977. NIOSH manual of analytical methods. 2nd ed. Vol 3. Washington, DC: U.S. Department of Health, Education, and Welfare, Public Health Service, Center for Disease Control, National Institute for Occupational Safety and Health, S274-1 - S274-7.

NIOSH. 1979. NIOSH manual of analytical methods. Vol. 5. Washington, D.C. U.S. Department of Health, Education, and Welfare, National Institute for Occupational Safety and Health. DHEW (NIOSH) Pub. No. 79-141.

NIOSH. 1984. NIOSH Manual of analytical methods, 3rd ed. Eller PM, ed. Cincinnati, OH: U.S. Department of Health and Human Services, National Institute for Occupational Safety and Health. Volume 1.

NIOSH. 1984a. NIOSH manual of analytical methods. Vol. 1: Method no. 5503. 3rd ed. Cincinnati, OH: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Institute for Occupational Safety and Health.

NIOSH. 1984b. NIOSH manual of analytical methods. Vol. 1: Method no. 8004. Cincinnati, OH: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Institute for Occupational Safety and Health.

NIOSH. 1989b. NIOSH manual of analytical methods. 3rd ed. Cincinnati, OPH: U.S. Department of Health and Human Services, Public Health Service, Center of Disease Control, National Institute for Occupational Safety and Health, 5510-I to 5510-4.

Noguchi T, Charman W, Stella V. 1985. Lymphatic appearance of DDT in thoracic or mesenteric lymph duct cannulated rats. *Int J Pharm* 24:185-192.

Norén K, Lunden A, Pettersson E, et al. 1996. Methylsulfonyl metabolites of PCBs and DDE in human milk in Sweden, 1972-1992. *Environ Health Perspect* 104(7):766-772.

Noren K, Sjoevall J. 1987. Analysis of organochlorine pesticides, polychlorinated biphenyls, dibenzo-p-dioxins and dibenzofurans in human milk by extraction with the lipophilic gel Lipidex® Noren K. J Chromator 422:103-115.

Norstrom RJ, Hallet DJ, Onuska FI, et al. 1980b. Mirex and its degradation products in Great Lakes herring gulls. Environ Sci Technol 14(7):860-866.

Norstrom RJ, Won HT, Holdrinet MVH, et al. 1980a. Gas-liquid chromatographic determination of mirex and photomirex in the presence of polychlorinated biphenyls: Interlaboratory study. J Assoc Off Anal Chem 63(1):37-42.

Nygren M, Hansson M, Sjöström M, et al. 1988. Development and validation of a method for determination of PCDDs and PCDFs in human blood plasma: A multivariate comparison of blood and adipose tissue levels between Vietnam veterans and matched controls. Chemosphere 17:1663-1692.

Oehme M, Mano S, Mikalsen A, et al. 1986. Quantitative Method for the determination of femtogram amounts of polychlorinated dibenzo-p-dioxins and dibenzofurans in outdoor air.

Ojala M. 1993. Simultaneous separation and determination of chlorobenzenes, PCBs, and chlorophenols using silica gel fractionation and GC-ECD analysis. Journal of High Resolution Chromatography 16: 679-682.

O'Keefe P, Meyer C, Smith R, et al. 1986. Reverse-phase adsorbent cartridge for trapping dioxins in drinking water. Chemosphere 15:1127-1134.

Oliver BG, Nicol KD. 1982b. Gas chromatographic determination of chlorobenzenes and other chlorinated hydrocarbons in environmental samples using fused silica capillary columns. Chromatographia 16:336-340.

Onuska FI, Comba ME, Coburn JA. 1980. Quantitative determination of mirex and its degradation products by high-resolution capillary gas chromatography/mass spectrometry. Anal Chem 52(14):2272-2275.

Onuska FI, Terry KA. 1993. Extraction of pesticides from sediments using a microwave technique. Chromatographia 36:101-104.

Onuska FL, Terry KA, Seech A, et al. 1994. Determination of toxaphene in soil by electron-capture negative-ion mass spectrometry and capillary column gas chromatography. J Chromatogr A 665: 125-132.

Opelario L, Rack E, Blotcky A, et al. 1983. Determination of chlorinated pesticides in urine by molecular neutron activation analysis. Anal Chem 55:677-681.

OSW. 1986a. Method 8081B - Organochlorine pesticides by gas chromatography. In: Test Methods for Evaluating Solid Waste, Physical/Chemical Methods SW 846. Office of Solid Waste, U.S. Environmental Protection Agency.

OSW. 1986b. Method 8250A - Semivolatile organic compounds by gas chromatography mass spectrometry (GC/MS). In: Test Methods for Evaluating Solid Waste, Physical/Chemical Methods SW 846. Office of Solid Waste, U.S. Environmental Protection Agency.

Patterson DG Jr, Fürst P, Alexander LR, et al. 1989b. Analysis of human serum for PCDDs/PCDFs: A comparison of three extraction procedures. *Chemosphere* 19:89-96.

Patterson DG Jr, Hampton L, Lapeza Cr Jr, et al. 1987a. High-resolution gas chromatographic/highresolution mass spectrometric analysis of human serum on a whole-weight and lipid basis for 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Anal Chem* 59:2000-2005.

Patterson DG Jr., Lapeza CR, Barnhart ER, et al. 1989. Gas chromatographic/mass spectrometric analysis of human serum for non-ortho (coplanar) and ortho substituted polychlorinated biphenyls using isotope-dilution mass spectrometry. *Chemosphere* 19:127-134.

Patterson DG Jr., Todd GD, Turner WE, et al. 1994. Levels of non-ortho-substituted (coplanar), monoand di-ortho-substituted polychlorinated biphenyls, dibenzo-p-dioxins, and dibenzofurans in human serum and adipose tissue. *Environ Health Perspect Suppl* 102(1):195-204.

Patterson DG, Hampton L, Lapeza CR, et al. 1987. High resolution gas chromatographic/highresolution mass spectrometric analysis of human serum on a whole weight and lipid basis for 2,3,7,8- tetrachlorodibenzo- p-dioxins. *Anal Chem* 59:2000-2005.

Pereira WE, Rostad CE, Sisak ME. 1985. Geochemical investigation of polychlorinated dibenzo-p-dioxins in the subsurface environment at an abandoned wood-treatment facility. *Environ Toxicol Chem* 4:629-639.

Petrick G, Schulz DE, Duinker JC. 1988. Clean-up of environment samples by high-performance liquid chromatography for analysis of organochlorine compounds by gas chromatography with electron-capture detection. *J Chromatogr* 435:241-248.

Polishuk ZW, Ron M, Wassermann M, et al. 1977a. Organochlorine compounds in mother and fetus during labor. *Environ Res* 13:278-294.

Polishuk ZW, Ron M, Wassermann M, et al. 1977b. Pesticides in people: Organochlorine compounds in human blood plasma and milk. *Pestic Monit J* 10(4):121-129.

Poon K-F, Lam PKS, Lam MHW. 1999. Determination of polychlorinated biphenyls in human blood serum by SPME. *Chemosphere* 39(6):905-912.

Prapamontol T, Stevenson D. 1991. Rapid method for the determination of organochlorine pesticides in milk. *J Chromatog* 552:249-257.

Que Hee S, Ward JA, Tabor MW, et al. 1983. Screening method for Aroclor 1254 in whole blood. *Anal Chem* 55: 157-160.

Radomski JL, Deichmann WB, Clizer EE, et al. 1968. Pesticide concentrations in the liver, brain, and adipose tissue of terminal hospital patients. *Food Cosmet Toxicol* 6:209-220.

Rahman MS, Bowadt S, Larsen B. 1993. Dual-column GC analysis of Mediterranean fish for ten organochlorine pesticides and sixty two chlorobiphenyls. *Journal of High Resolution Chromatography* 16:731-735.

Rappe C, Kjeller LO, Andersson R. 1989c. Analyses of PCDDs and PCDFs in sludge and water samples. *Chemosphere* 19:13-20.

Rappe C, Kjeller LO, Bruckmann P, et al. 1988. Identification and quantification of PCDDs and PCDFs in urban air. *Chemosphere* 17:3-20.

Reding R. 1987. Chromatographic monitoring methods for organic contaminants under the Safe Drinking Water Act. *J Chromatogr Sci* 25:338-344.

Richardson A, Robinson J. 1971. The identification of a major metabolite of HEOD (dieldrin) in human feces. *Xenobiotica* 1(3):213-219.

Ritcey WR, Savary G, McCully KA. 1972. Organochlorine insecticide residues in human milk, evaporated milk and some milk substitutes in Canada. *Can J Pub1 Hlth* 63: 125-132.

Roberts JW, Camann DE. 1989. Pilot study of a cotton glove press test for assessing exposure to pesticides in house dust. *Bull Environ Contam Toxicol* 43:717-724.

Rohrig L, Meisch H. 2000. Application of solid phase micro extraction for the rapid analysis of chlorinated organics in breast milk. *Fresenius J Anal Chem* 366:106-111.

Russo MV, Goretti G, Liberti A. 1993. Rapid determination of chlorinated pesticides using CN-bonded cartridges followed by GC-ECD. *Chromatographia* 35:290-294.

Ryan JJ, Lizotte R, Lewis D. 1987a. Human tissue levels of PCDDs and PCDFs from a fatal pentachlorophenol poisoning. *Chemosphere* 16:1989-1996.

Ryan JJ, Lizotte R, Lewis D. 1987b. Human tissue levels of PCDDs and PCDFs from a fatal pentachlorophenol poisoning, *Chemosphere* 16: 1989-1996.

Ryan JJ, Schecter A, Sun NF, et al. 1986. Distribution of chlorinated dibenzo- p-dioxins and chlorinated dibenzo-furans in human tissues from the general population. In: Rappe C, Choudhury, G,

Saady JJ, Fitzgerald RL, Poklis A. 1992. Measurement of DDT and DDE from serum, fat, and stool specimens. *J Environ Sci Health A27(4)*:967-981.

Saito I, Kawamura N, Uno K, et al. 1985. Determination of chlordane in human blood by gas chromatography. *Analyst* 110:263-267.

Saleh MA, Casida JE. 1977. Consistency of toxaphene composition analyzed by open tubular column gas-liquid chromatography. *J Agric Food Chem* 25:63-68.

Sasaki K, Ishizaka T, Suzuki T, et al. 1991b. Organochlorine chemicals in skin lipids as an index of their accumulation in the human body. *Arch Environ Contam Toxicol* 21:190-194.

Schecter A, Tiernan T. 1985. Occupational exposure to polychlorinated dioxins, polychlorinated furans, polychlorinated biphenyls, and biphenylenes after an electrical panel and transformer accident in an office building in Binghamton, NY. *Environ Health Perspect* 60:305-313.

Schmidt LJ, Hesselberg RJ. 1992. A mass spectroscopic method for analysis of AHH-inducing and other polychlorinated biphenyl congeners and selected pesticides in fish. *Arch Environ Contam Toxicol* 23(1):37-44.

Schuman RF, Hunter KW. 1988. Ultra sensitive bioassay for dioxin. Electric Power Research Institute Report, 7-11 - 7-14.

Schwartz TR, Tillitt DE, Feltz KP, et al. 1993. Determination of mono- and non-O,O'-chlorine substituted polychlorinated biphenyls in Aroclors and environmental samples. *Chemosphere* 26(8):1443-1460.

Seiber JN, Landrum PF, Madden SC, et al. 1975. Isolation and gas chromatographic characterization of some toxaphene components. *J Chromatogr* 114:361-368.

Seidel V, Linder W. 1993. Universal sample enrichment technique for organochlorine pesticides in environmental and biological samples using a redesigned simultaneous steam distillation- solvent extraction apparatus. *Anal Chem* 65:3677-3683.

Shan TH, Hopple JA, Foster GD. 1994. Alternative tissue analysis method developed for organochlorine contaminants in aquatic organisms. *Bull Environ Contam Toxicol* 53:382-389.

Simon N, Guzzetta J, Thielen D. 1989. GC/MS and GC/MS/MS: Complimentary techniques for the quantitation of tetrachlorodibenzo-p-dioxin in soils and sediments from industrial sites. *Chemosphere* 18:163-168.

Singh J, Miles W, Barrette JP. 1985. Determination of polychlorinated dibenzo-p-dioxins in technical pentachlorophenol. *J Assoc Off Anal Chem* 68:583-586.

Smith LM, Stalling DH, Johnson JL. 1984. Determination of part-per-trillion levels of polychlorinated dibenzofurans and dioxins in environmental samples. *Anal Chem* 56: 1830- 1842.

Smith RM, O'Keefe PW, Hilker DR, et al. 1986b. Sampling, analytical method and results for chlorinated dibenzo-p-dioxins and chlorinated dibenzofurans from incinerator stack effluent and contaminated building indoor air samples. In: Rappe C, Choudhary G, Keith LH, eds. *Chlorinated dioxins and dibenzofurans in perspective*. Chelsea, MI: Lewis Publishers, Inc., 93-108.

Smrek AL, Needham LL. 1982. Simplified cleanup procedures for adipose tissue containing polychlorinated biphenyls, DDT, and DDT metabolites. *Bull Environ Contam Toxicol* 28:718-722.

Snyder JL, Grob RL, McNally ME, et al. 1992. Comparison of supercritical fluid extraction with classical sonication and soxhlet extractions for selected pesticides. *Anal Chem* 64:1940-1946.

Sperling KR, Janssen D, Wulf T. 1985. Gas chromatographic-mass spectrometric investigations on lipophilic anthropogenic substances in marine organisms. I. A clean-up procedure consisting of liquid-liquid and reversed phase (octadecyl)-partitioning. *Fresenius' Journal of Analytical Chemistry* 320:1-5.

Stacey CI, Tatum T. 1985. House treatment with organochlorine pesticides and their level in human milk--Perth, Western Australia. *Bull Environ Contam Toxicol* 35:202-208.

Stachel B, Dougherty RC, Lahl U, et al. 1989. Toxic environmental chemicals in human semen: Analytical method and case studies. *Andrologia* 21:282-291.

Stalling DL, Peterman PH, Smith LM, et al. 1986. Use of pattern recognition in the evaluation of PCDD and PCDF residue data from GC/MS analysis. *Chemosphere* 15:1435-1443.

Stanker LH, Watkins B, Rogers N, et al. 1987. Monoclonal antibodies for dioxin: Antibody characterization and assay development. *Toxicology* 45:229-243.

Stanley JS, Boggess KE, Onstott J, et al. 1986. PCDDs and PCDFs in human adipose tissue from the EPA FY82 NHATS repository. *Chemosphere* 15:1605-1612.

Stanley JS. 1986. Broad scan analysis of the FY82 National human adipose tissue survey specimens: Volume 1. Executive summary. Washington, D.C.: U.S. Environmental Protection Agency. EPA 560/5-86-035.

Swackhamer DL, Armstrong DE. 1987. Distribution and characterization of PCBs in Lake Michigan water. *J Great Lakes Res* 13:24-36.

Swackhamer DL, Charles MJ, Hites RA. 1987. Quantitation of toxaphene in environmental samples using negative ion chemical ionization mass spectrometry. *Anal Chem* 59:913-917.

Takizawa Y, Muto H. 1987. PCDDs and PCDFs carried to the human body from the diet. *Chemosphere* 16:1971-1975.

Tang PH, Ho JS, Eichelberger JW. 1993. Determination of organic pollutants in reagent water by liquid-solid extraction followed by supercritical fluid elution. *J AOAC International* 76:72-82.

Tashiro C, Clement RE, Reid N, et al. 1989. Determination of dioxins and furans in precipitation collected in urban and rural Ontario Canada locations. *Chemosphere* 19:535-540.

Tashiro S, Matsumura F. 1978. Metabolism of trans-nonachlor and related chlordane components in rat and man. *Arch Environ Contam Toxicol* 7:113-127.

Tewari SN, Sharma IC. 1977. Isolation and determination of chlorinated organic pesticides by thin-layer chromatography and the application to toxicological analysis. *J Chromatogr* 131:275-284.

Thier H-P, Zeumer H. 1987a. Organochlorine, organophosphorus nitrogen-containing and other pesticides. Deutsche Forschungsgemeinschaft: Manual of pesticide residue analysis. Vol. 1. Weinheim, Germany: VCH, 383-400.

Thier H-P, Zeumer H. 1987b. Organochlorine and organophosphorus pesticides. Deutsche Forschungsgemeinschaft: Manual of pesticide residue analysis. Vol. 1. Weinheim, Germany. VCH, 298-307.

Thomas TC, Nishioka YA. 1985. Sampling and analytical procedures for evaluation of airborne chlordane levels in family quarters. *Environmental Toxicology and Chemistry* 4:143-147.

Tiernan TO, Solch JG, Garrett JG, et al. 1990. A concerted analytical method for determination of various halogenated and related bioaccumulating compounds in fish and sediments. *Brominated Compound, Short-Chain Aliphatics* 2:225-228.

Tiernan TO, Taylor ML, VanNess GF, et al. 1984. Analyses of human tissues for chlorinated dibenzo- p-dioxins and chlorinated dibenzofurans: The state of the art. In: Public health risks of the dioxins. Lowrance, WW, ed. New York, NY: Rockefeller University, 31-56.

Tiernan TO, Wage1 DJ, Garrett JH, et al. 1989. Laboratory and field tests to demonstrate the efficacy of KPEG reagent for detoxification of hazardous wastes containing polychlorinated dibenzop-dioxins (PCDD) and dibenzofurans (PCDF) and soils contaminated with such chemical wastes. *Chemosphere* 18:835-841.

Tojo Y, Wariishi M, Suzuki Y, et al. 1986. Quantitation of chlordane residues in mothers' milk. *Arch Environ Contam Toxicol* 15:327-332.

Tomkins BA, Merriweather R, Jenkins RA. 1992. Determination of eight organochlorine pesticides at low nanogram/liter concentrations in groundwater using filter disk extraction and gas chromatography. *J AOAC International* 75:1091-1099.

Tondeur Y, Beckert WF, Billels S, et al. 1989. Method 8290: An analytical protocol for the multimedia characterization of polychlorinated dibenzodioxins and dibenzofurans by high-resolution gas chromatography/high-resolution mass spectrometry. *Chemosphere* 18: 119- 131.

Tong HY, Shore DL, Karasek FW, et al. 1984. Identification of organic compounds obtained from incineration of municipal waste by high-performance liquid chromatographic fractionation and gas chromatography-mass spectrometry. *J Chromatogr* 285:423-441.

Trotter WJ, Dickerson R. 1993. Pesticide residues in composited milk collected through the U.S. Pasteurized Milk Network. *J AOAC Int* 76(6):1220-1225.

Tulp MTM, Hutzinger O. 1978. Rat metabolism of polychlorinated dibenzo-p-dioxins. *Chemosphere* 7:761-768.

Van den Berg M, Heeremans C, Veenhoven E, et al. 1987b. Transfer of polychlorinated dibenzo-p-dioxins and dibenzofurans to fetal and neonatal rats. *Fundam Appl Toxicol* 9:635-644.

Van den Berg M, Van der Wielen FWM, Olie K, et al. 1986b. Presence of PCDDs and PCDFs in human breast milk from the Netherlands. *Chemosphere* 15:693-706.

Van den Berg M, Van Wijnen J, Wever H, et al. 1989. Selective retention of toxic polychlorinated dibenzo-p-dioxins and dibenzofurans in the liver of the rat after intravenous administration of a mixture. *Toxicology* 55: 173-182.

Van der Velde EG, Marsman JA, De Jong AP, et al. 1994. Analysis and occurrence of toxic planar PCBs, PCDDs and PCDFs in milk by use of carbosphere activated carbon. *Chemosphere* 28(4):693-702.

Vaz R, Blomkvist G. 1985. Traces of toxaphene components in Swedish breast milk analyzed by capillary GC using ECD, electron impact and negative ion chemical ionization MS. *Chemosphere* 14:223-231.

Verma A, Pillai MKK. 1991a. Bioavailability of soil-bound residues of DDT and HCH to certain plants. *Soil Biol Biochem* 23(4):347-351.

Wage1 DJ, Tiernan TO, Taylor ML, et al. 1989. Assessments of ambient air sampling techniques for collecting airbourne polyhalogenated dibenzo- p-dioxins (PCDDs) dibenzofurans (PCDFs) and biphenyls (PCBs). *Chemosphere* 18:177-184.

Wagner SL, Durand LR, Inman RD, et al. 1991. Residues of pentachlorophenol and other chlorinated contaminants in human tissues: Analysis by electron capture gas chromatography and electron capture negative ion mass spectrometry. *Arch Environ Contam Toxicol* 21(4):596-606.

Waliszewski S, Szymczynski G. 1983. Determination of selected chlorinated pesticides, bound and free, in human semen. *Arch Environ Contam Toxicol* 12:577-580.

Waliszewski SM, Szymczynski GA. 1985. Inexpensive, precise method for the determination of chlorinated pesticide residues in soil. *J Chromatogr* 321:480-483.

Walker AIT, Stevenson DE, Robinson J, et al. 1969. The toxicology and pharmacodynamics of dieldrin (HEOD): Two-year oral exposures of rats and dogs. *Toxicol Appl Pharmacol* 15:345-373.

Wallace BG, Sherren AJ. 1986. Measurement of aldrin by gas chromatography with sorbent-filled trapping tube. *J AOAC* 69(1):93-97.

Walters SM. 1986. Cleanup of samples. In: Zweig G, Sherma J. eds. *Analytical methods for pesticides and plant growth regulators*, Vol 15, Chap. 3. New York, NY: Academic Press:67-110.

Wariishi M, Nishiyama K. 1989. Observations on the progress of chlordane contamination in humans by blood and sebum analysis. *Arch Environ Contam Toxicol* 18:501-507.

Wariishi M, Suzuki Y, Nishiyama K. 1986. Chlordane residues in normal human blood. *Bull Environ Contam Toxicol* 36:635-643.

Welsh MS. 1995. Extraction and gas chromatography/electron capture analysis of polychlorinated biphenyls in railcar paint scrapings. *Appl Occup Environ Hyg* 10(3):175-181.

Wendling JM, Orth RG, Poiger H. 1990. Determination of [3H]-2,3,7,8-tetrachlorodibenzo-p-dioxin in human feces to ascertain its relative metabolism in man. *Anal Chem* 62:796-800.

WHO. 1984. World Health Organization. Camphechlor. Environ Health Criteria 45: 1-66.

Wigfield YY, Grant R. 1992. Evaluation of an immunoassay kit for the detection of certain organochlorine (cyclo-diene) pesticide residues in apple, tomato, and lettuce. Bull Environ Contam Toxicol 49:342-347.

Williams DT, LeBel GL, Junkins E. 1988a. Organohalogen residues in human adipose autopsy samples from six Ontario municipalities. J Assoc Off Anal Chem 71:410-414.

Williams S. 1984. Official methods of analysis. Association of Official Analytical Chemists, Inc., Arlington, VA.

Wirth EF, Chandler GT, Dipinto LM, et al. 1994. Assay of polychlorinated biphenyl bioaccumulation from sediments by marine benthic copepods using a novel microextraction technique. Environ Sci Technol 28:1609-1614.

Yagüe C, Bayarri S, Lázaro R, et al. 2001. Multiresidue determination of organochlorine pesticides and polychlorinated biphenyls in milk by gas chromatography with electron-capture detection after extraction by matrix solid-phase dispersion. J AOAC Int 84:1561-1568.

Young S, Clower Jr M, Roach JAG. 1984. Method for determination of organohalogen pesticide residues in vegetable oil refinery by-products. J Assoc Off Anal Chem 67:95-106.

Zacharewski T, Safe L, Safe S, et al. 1989. Comparative analysis of polychlorinated dibenzop-dioxin and dibenzofuran congeners in Great Lakes fish extracts by gas chromatography-mass spectrometry and in vitro enzyme induction activities. Environmental Science and Technology 23:730-735.

Zacharewski T, Safe L, Safe S, et al. 1989. Comparative analysis of polychlorinated dibenzo-p-dioxin and dibenzofuran congeners in Great Lakes fish extracts by gas chromatography and in vitro enzyme induction activities. Environ Sci Technol 23:730-735.

Zupancic-Kralj L, Jan J, Marsel J. 1992. Assessment of polychlorobiphenyls in human/poultry fat and in hair/plumage from a contaminated area. Chemosphere 25(12):1861-1867.